

UNIVERSITI TEKNOLOGI MARA

**BIOMARKERS OF ENDOTHELIAL
ACTIVATION AND ATP-BINDING
CASSETTE TRANSPORTER A1
(*ABCA1*) GENE VARIANTS IN
SUBJECTS WITH LOW HDL-C
CONCENTRATION**

WAN NOR HANIS BINTI WAN AHMAD

Thesis is submitted in fulfillment
of the requirements for the degree of
Master of Science


Faculty of Medicine

December 2015

AUTHOR’S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of student	:	Wan Nor Hanis Binti Wan Ahmad
Student I.D. No.	:	2011436178
Programme	:	Master of Science (Medicine) – MD780
Faculty	:	Faculty of Medicine
Thesis Title	:	Biomarkers of Endothelial Activation and ATP-Binding Cassette Transporter A1 (<i>ABCA1</i>) Gene Variants in Subjects with Low HDL-c Concentration
Signature of Student	:	
Date	:	December 2015

ABSTRACT

Soluble vascular cell adhesion molecule-1 (sVCAM-1), intercellular cell adhesion molecule 1 (sICAM-1), and E-selectin are biomarkers reflecting endothelial activation (EA), a key event in atherosclerosis. High-density lipoprotein cholesterol (HDL-c) is protective against atherosclerosis by reverse cholesterol transport (RCT) but its association with EA is not well established. ATP-binding Cassette Transporter A1 (*ABCA1*) is the main transport protein in RCT and its gene variants have been linked to low HDL-c concentration. Thus, this study aimed to (a) compare the concentration of biomarkers of EA between low HDL-c subjects and normal controls (NC), (b) examine the correlation and association between HDL-c and biomarkers of EA, (c) investigate whether HDL-c concentration is an independent predictor for these biomarkers, and (d) determine the genetic variants of *ABCA1*. A total of 207 subjects with low HDL-c and 215 age-, gender-, ethnic-, smoking-, diabetic- and blood pressure-matched NC were recruited. Fasting blood samples were collected for biomarkers of EA and DNA extraction. Targeted regions of *ABCA1* gene were amplified and sequenced. The detection of the SNPs were analysed using BioEdit. Subjects with low HDL-c had greater concentrations EA biomarkers compared to controls. There were significant negative correlations and association between HDL-c concentration and EA biomarkers. HDL-c was an independent predictor for sVCAM-1. DNA sequencing of *ABCA1* revealed six genetic variants and two SNPs were found to be significantly different between the low HDL-c subjects and NC in both genotype and allele frequencies. In conclusion, low serum HDL-c concentration is strongly correlated and associated with enhanced status of EA which strongly supports its role in the pathogenesis of atherosclerosis. The findings of genetic variation in *ABCA1* suggest that this gene may play an important role in HDL deficiency amongst Malaysians.

ACKNOWLEDGEMENT

Firstly, I am thankful to Allah for His blessing and giving me the opportunity to embark on my Master Programme by completing this winding and challenging journey successfully. In this short page, I would like to convey my sincere gratitude and deepest appreciation to a number of people, as without them, this thesis would not be possible. To my great mentor and supervisor Prof Dr Hapizah Mohd Nawawi, I am really grateful for her encouragement, supports and precious guidance during this whole arrangement of research and the preparation of thesis. Also not to forget to my co-supervisors, Dr Thuhairah Abdul Rahman and Dr Hoh Boon Peng, as they have always given me full support, professional assistance and useful suggestions throughout the laboratory works and thesis write-up.

My deepest gratitude to staff members of Center for Pathology and Diagnostic Research Laboratory (CPDRL) and Institute of Medical Molecular Biotechnology (IMMB), Universiti Teknologi MARA (UiTM) for their willingness to share expertise in both biochemical assays and molecular work as well as allowing me to work in their laboratory until the end of the experiments. This also includes, to co-researchers of the research and colleagues for giving their support either directly and indirectly. My heartfelt thanks goes to my colleagues, Miss Nurul Atiqah Mohd Mokhsin and Miss Farah Hanis Sakri as we have shared many memories together in doing our research by exchanging our thoughts, guidance and giving support to each other from the beginning until the end of this research.

Apart from that, I would like to take this opportunity to extend my thanks to the Ministry of Higher Education, Long Term Research Grant (LRGS) for the research grant awarded to my supervisor, Faculty of Medicine and Institute of Postgraduate Studies (IPSiS), UiTM.

Finally, my greatest appreciation to my family members especially to my father, Mr Wan Ahmad Abdullah, my husband, Mr Muhammad Alqaf Azmi and my sister, Mrs Wan Nor Aliza Wan Ahmad for their full encouragement, financial support, motivation, prayers and being understandable with my research. I am also dedicating this thesis in remembrance of my very dear late mother, Mrs Siti Hawa Abu Bakar for her words which inspire me to keep on learning and never give up in life. Without all of them, I could not have completed this whole thing successfully.

In conclusion, throughout this research I have learnt a lot of things. I have made many incredible findings and good friends, interacting with great expertise and professional persons, working under stress and not to mention, dealing with problems efficiently. All of these had made me realized that research is a very fascinating and challenging field which full of surprises. If given a chance, I would love to continue in this field by giving contribution to the society as well as mankind.

CHAPTER ONE

INTRODUCTION

1.1 RESEARCH BACKGROUND

Approximately \$110 billion has been spent in the year 2010 alone for the cost of health care services, medications and loss of productivity in the US due to coronary artery disease (CAD), and by 2020 (Chockalingam et al., 2000) it is estimated that CAD will be the number one cause of disability and death worldwide which is nearly 25 million of death (Murray & Lopez, 1997). According to The National Health and Morbidity Survey in 2011, Malaysia was reported as having a high prevalence of chronic disease including high cholesterol. In 2010, about 40% of death in Malaysia was estimated to be caused by chronic diseases. Whilst, data from World of Health Organization published in 2014, CHD death in Malaysia reached 29,363 or 23.10% of total death in which the death rate is 150.11 per 100,000 of population.

To date, CAD remains one of the major causes of death in the developed and developing countries (Xu et al., 2013) and is commonly due to atherosclerosis, a systemic and complex disease in which lipid and fibrous material accumulate within the intimal layer of large to medium-sized arteries. It is also the most important and common underlying cause of myocardial infarction, ischaemic stroke, and peripheral vascular diseases apart from being the major cause of chronic heart failure and vascular dementia. It can start as early as foetal life (Palinski & Napoli, 2002), progressing slowly throughout childhood and adolescence and accelerating in adult life. It also involves indolent inflammation, endothelial activation, oxidative stress and prothrombogenesis (Virmani, Kolodgie, Burke, Farb, & Schwartz, 2000).

Inflammation, endothelial activation, oxidative stress and prothrombogenesis are significant contributors to the development and progression of atherosclerosis. Several biomarkers have been established to reflect the status of these processes. Biomarkers such as high sensitivity C-reactive protein (hsCRP) and Interleukin-6 (IL-6) are indicators of inflammation (Schuett, Luchtefeld, Grothusen, Grote, & Schieffer, 2009), while endothelial activation is reflected through the various adhesion molecules such as soluble vascular cell adhesion molecule-1 (sVCAM-1), soluble